

**Bruce R. Maddern M.D., P.A.**  
**Post Frenuloplasty (procedure to resolve “tongue-tie”) Instructions**

Ankyloglossia (“tongue-tie”) is a condition present from birth in which movement of the tongue tip is restricted due to an unusually short and/or thick frenulum. A frenulum is a band of tissue that anchors the tongue to the floor of the mouth. When the frenulum is too short or tight, the tongue tip is unable to move freely. This may result in difficulty with early bottle or breast-feeding or speech articulation.

The most common reason to perform frenuloplasty in a newborn is difficulty with feeding or latching. It is possible to perform the frenuloplasty in office for infants during the first few months of life. The baby is able to feed immediately following the procedure. In infants 6 months old or younger, frenuloplasty can normally be performed in the office with a local anesthetic.

In toddlers it may be necessary to perform a frenuloplasty because of speech articulation issues. If a frenuloplasty is required for this age group the procedure occurs in the operating room under general anesthesia. The entire procedure usually is complete within 15 minutes. Speech therapy is necessary to ensure improved mobility of the tongue for older children. General risks from frenuloplasty include bleeding, pain, scarring, and infection.

**General Information**

- The patient may resume normal activity following surgery, including normal feeding schedule (bottle, breast, or solid food).
- The patient may return to daycare the next day.
- Some discomfort is expected following this procedure. You may treat your child’s pain with Tylenol or ibuprofen.
- Ice pops are a good way to numb the area. After the child finishes the ice pop, use the stick to stretch out the area under the tongue at least once a day to prevent scarring.
- Following the procedure it is normal to see a dissolvable stitch or an area under the tongue that may be yellow, white, brown, or gray. This is a part of the normal healing process.
- It is normal to have an increase in drooling the first few days following the procedure. This will resolve on its own.
- Toddlers and older children who undergo frenuloplasty for speech articulation issues must be enrolled in speech therapy and can resume sessions in 1-2 weeks.
- Please make sure to follow up with our office 2 weeks after the procedure.

Updated 2/10

Bruce R. Maddern, M.D., P.A.  
A division of Florida Pediatric Associates, LLC  
10475 Centurion Parkway N. Ste. 302  
Jacksonville, FL 32256  
[www.entforkids.com](http://www.entforkids.com)  
(904) 398-5437

