




Adenoidectomy

Post-operative instructions



Dr Andrew Simonsen, DO
Dr Angela Black, MD
Dr Steven Andreoli, MD

Adenoidectomy (Removal of the adenoids) The adenoids are located behind the nose and hidden from view by the palate (roof of the mouth). Chronic obstructive rhinitis, frequent ear infections, nasal airway obstruction, or obstruction of the eustachian tube (a passage that connects the middle ear to the back of the throat) are some reasons for removal (adenoidectomy).

Surgery is performed under general anesthesia, for most kids this is an outpatient surgery (This means you have surgery and go home the same day). If your child has multiple medical problems or severe obstructive sleep, they may need to stay in the hospital one night for observation.

General Instructions

Following surgery most kids will recover within 1-3 days. Some children will complain of a sore throat for a few days after surgery. Snoring and mouth breathing is normal after surgery due to the swelling in the back of the nose.

Drinking fluids and nutrition are very important to ensure healing. Your child may eat a normal diet after surgery. There are no diet restrictions.

Antibiotics are not typically needed after surgery. Some children may have a low-grade fever, this is normal for a few days after surgery. Some kids can have bad breath for a period after surgery, this is normal.

Normal activities may be resumed as soon as your child feels up to it. Most children can go back to school/daycare the next day after surgery.

Who to call?

Please direct all questions and concerns related to your child's surgery to our office (904) 398-5437

Pain Control

A mild amount of throat discomfort is to be expected after adenoidectomy. Some children will complain of headaches or ear pain for a few days after surgery, this is usually relieved with pain medication.

Children should be given Acetaminophen (Tylenol®) and Ibuprofen (Advil® or Motrin®) alternating every 3 hours after surgery.

CAUTION: Medications containing Ibuprofen should not be taken during the 10 days PRIOR to surgery. Medications containing Codeine or Toradol should NOT be used for children after adenoidectomy.

The Operative Site

A foul odor is common from the mouth/nose. Teeth brushing is allowed and encouraged. Using a room humidifier (at night) or nasal saline spray is recommended to help with oral dryness, snoring and congestion.

Post-Operative Pain Medication Dosing Guide

Child's Weight	Daily minimal fluid intake	Tylenol (Acetaminophen) 160mg/5ml	Motrin/ Advil (Ibuprofen) 100mg/5ml	Hycet Dose (Based on specific weight)
	Ounces (oz)	Every 4-6 hours as needed for pain Max: 5 doses in 24 hours	Every 6-8 hours as needed for pain Max: 4 doses in 24 hours	Hycet contains Acetaminophen so do not give both at the same time.
<17 lbs (5-8 kg)	35 oz	½ tsp (2.5 ml)	½ tsp (2.5ml)	
17-20 lbs (8-9 kg)	45 oz	¾ tsp (3.7ml)	¾ tsp (3.7ml)	
20-24 lbs (9-11 kg)	55 oz	¾ tsp (3.7ml)	1 tsp (5 ml)	
24-35 (11-16 kg)	65 oz	1 tsp (5 ml)	1½ tsp (7.5 ml)	
35-50 lbs (16-23 kg)	>65 oz	1½ tsp (7.5 ml)	2 ml (10 ml)	
50-65 lbs (23-29 kg)	>65 oz	2 ml (10 ml)	2 ½ tsp (12.5 ml)	
65-80 lbs (29-37 kg)	>65 oz	3 tsp (15 ml)	3 ½ tsp (17.5 ml)	
89-95 lbs (37-43 kg)	>65 oz	3 ½ tsp (17.5 ml)	4 tsp (20 ml)	
>95 lbs (>43 kg)	>65 oz	4 tsp (20 ml)	4 tsp (20 ml)	

IMPORTANT: Do not give Acetaminophen and Hycet simultaneously. Hycet contains added acetaminophen and combining the two may lead to toxic medication levels.

IMPORTANT: There is no substitute for Hydrocodone (Lortab or Hycet) and should only be used if Tylenol/Advil fail.

Tylenol (Acetaminophen) is available as a suppository (you can get this at any pharmacy without a prescription) and may be recommended if your child is having difficulty swallowing.

Liquid/syrup medications may be mixed with a clear “tasty” liquid to make it easier to swallow.

Benadryl with Advil/Motrin is a good choice for additional pain relief if Tylenol/Advil alone are not sufficient.

Benadryl OTC (Diphenhydramine) – 12.5mg/5ml, May be given up to every 6 hours

2-6 yrs old: 2.5ml

6-12 yrs old: 5ml

>12 yrs old: 10ml

Make a Medication Record to write down the time and medication you are giving. Especially if you are giving medication for days this will help you remember when you gave the last medication, so you do not miss a dose.

Updated 11/2022 AB
A Division of Florida Pediatric Associates, LLP
10475 Centurion Parkway N., Suite 302, Jacksonville, Florida 32256
904-398-KIDS (5437) FAX 904-398-3077
WWW.ENTFORKIDS.COM